

This information sheet is intended for health professional use. A consumer version will be available from the IMAC COVID-19 Education website soon.

COVID-19 and COVID-19 vaccination in pregnancy

Pregnant women/people are routinely offered the COVID-19 vaccine, Comirnaty, which is recommended at any stage of pregnancy.

COVID-19 vaccination with the Comirnaty™ mRNA COVID-19 vaccine (Pfizer/BioNTech) is available to anyone from 12 years of age. Two doses are given at least 21 days apart.

Internationally, many thousands of people have been given this vaccine while pregnant and large-scale surveillance data (presented below) indicate that there are no safety concerns with administering this COVID-19 vaccine in any stage of pregnancy. Vaccinating during pregnancy may also offer temporary protection for newborns via passive transfer of antibody across the placenta and in breastmilk.

Anyone with questions or concerns about receiving this vaccine in pregnancy is advised to discuss these with their health professional. Everyone has a right to make an informed decision about receiving the vaccine.

Routine pregnancy testing before COVID-19 vaccination is not required and for those who are planning pregnancy, it is not necessary to delay pregnancy after receiving a COVID-19 vaccine.

COVID-19 vaccination does not replace the need for simple measures that reduce the risk of disease transmission such as physical distancing, handwashing and use of appropriate personal protective equipment (PPE), as needed. It is also advisable for all household and health care contacts of a pregnant people, new mothers and their infants be fully vaccinated against COVID-19, as age appropriate.

When available, the use of the adenovirus vector COVID-19 Vaccine AstraZeneca in pregnant people/women needs to be based on an assessment of whether the benefits of vaccination with this vaccine outweigh the potential risks. Comirnaty (mRNA vaccine) is the preferred vaccine for use during pregnancy.

COVID-19 disease in New Zealand

Community transmission of COVID-19 is increasing in New Zealand, despite ongoing efforts to manage the international border, rapid identification and comprehensive contact tracing of community cases, which have slowed community spread of the disease.¹ High vaccination coverage is now needed to protect our communities, particularly those who are highest risk from COVID-19.

COVID-19 disease and pregnancy

Pregnant women/people are at a much higher risk from COVID-19 hospitalisation and intensive care unit admission than non-pregnant women.

Due to changes that take place during pregnancy, healthy pregnant women who are infected with the SARS-CoV-2 virus have an increased risk of severe COVID-19 disease compared with non-pregnant women.²

³ To prevent the mother's immune system from harming her growing baby, her immune system is

mRNA COVID-19 vaccine Pfizer/BioNTech [mRNA-CV]

For health professional use

temporarily suppressed making her more susceptible to infection⁴. In addition, as her baby grows, the mother's lung capacity decreases and the volume of blood being pumped around her body increases, requiring more oxygen and her heart to work harder.⁴⁻⁶ As a result, COVID-19 infection when pregnant or soon after pregnancy substantially increases the risk of severe disease and breathing difficulties.

Although the risk of catching COVID-19 is not increased, pregnant women with a COVID-19 diagnosis are over three times more likely to have severe disease and require hospital admission, and five times more likely to require intensive care unit admission than non-pregnant women.^{7, 8} The risk for severe disease is higher for women with an ethnic minority background. In the US, Asian and native Hawaiian/Pacific Islander women were disproportionately at great risk for ICU admission.³ The risks in pregnancy increases from the age of 35 years and for those who have a chronic condition, such as obesity, high blood pressure or pre-existing diabetes.⁹ During 2020, twice as many pregnant women with symptomatic COVID-19 died than those with COVID-19 who were not pregnant.^{3, 9}

According to data (peer-review pending) collected by the UK Obstetric Surveillance System, the proportion of pregnant women hospitalised with moderate or severe COVID-19 increased significantly as new variants of SARS-CoV-2 virus emerged (from 24%, to 36% and 45% with wildtype, Alpha and Delta variants, respectively). Those with Delta COVID-19 were at the highest risk of being hospitalised with pneumonia. Eleven pregnant women with COVID-19 died compared with one in the comparator group of women without COVID-19. No fully vaccinated pregnant women were admitted to hospital for COVID-19.¹⁰

COVID-19 increases the risk of complications in pregnancy, such as preeclampsia, hypertension and poor fetal growth.

A multinational cohort study found that women with COVID-19 infection who experienced fever and shortness of breath for any duration was associated with two and a half times increased risk for severe maternal complications and five times increased risk in neonatal clinical complications. The risks of medically indicated preterm birth and preeclampsia were twice as high in mothers with COVID-19.⁷

Although COVID-19 infection early in pregnancy is not associated with increased risk for pregnancy loss,² an increase in adverse pregnancy outcomes (gestational diabetes and hypertension, preeclampsia and poor fetal growth) has been observed during the COVID-19 pandemic in the US.¹¹ COVID-19 in women with underlying health conditions significantly increases the risk of preterm labour, preeclampsia and eclampsia than those without underlying conditions.¹²

COVID-19 increases the risk of preterm birth, caesarean section and baby admission to neonatal intensive care.

Babies are up to seven times more likely to be born preterm to mothers with COVID-19 and up to five times more likely to require neonatal intensive care when compared with babies born to mothers without the disease.⁹ Out of concern for the maternal health, the babies of women hospitalised with COVID-19 are more likely to be born by caesarean section and delivered early.¹³ As reported by the Ministry of Health, in 2018, 7.5% of babies were born preterm (before 37 weeks), in comparison, about half of the babies born to mothers with COVID-19 are being delivered early. The risk for preterm birth is higher among those mothers with underlying health conditions such as hypertension, diabetes or obesity.¹⁴ COVID-19 in pregnancy does not appear to affect the rates of neonatal death and stillbirth.¹³

mRNA COVID-19 vaccine Pfizer/BioNTech [mRNA-CV]

For health professional use

The risk of transmitting SARS-CoV-2 infection from the mother to her newborn appears to be small, is often asymptomatic and rarely severe, although the risk of severe disease in newborns is higher than for older infants.^{15, 16} One study in Italy found around half of the neonates infected with SARS-CoV-2 were likely to have been infected by family members or health care workers.² Note: to date in New Zealand, no women or their babies have been infected with COVID-19 by health workers.

How the COVID-19 vaccine works

Comirnaty COVID-19 vaccine contains messenger ribonucleic acid (mRNA) inside a lipid bubble which is delivered to muscle cells in the arm. It is not a live vaccine and contains no part of the SARS-CoV-2 virus. The mRNA and its protective bubble are so fragile that the vaccine needs to be stored at very cold temperatures to stop it from degrading too quickly.

After injection, the mRNA is taken into the cell where it delivers the instructions to make replicas of the protein (the spike protein) that the SARS-COV-2 virus uses to infect our cells. These cells use this protein to specifically activate the immune response against COVID-19 virus, in the same way as the more conventional vaccine antigens. The quantity of this protein produced after vaccination is much lower than the amount seen in people with COVID-19 infection with the virus spreading throughout their body. Furthermore, as soon as it is produced, this protein is dismantled inside these specialist cells and the pieces are used to activate the immune response.

The mRNA breaks down quickly after this process takes place. It is unable to enter the cell nucleus, cannot integrate with DNA,¹⁷ and will not cause genetic changes in the mother or the baby. The remainder of the vaccine components are also rapidly cleared from the body within a few days after vaccination, as for other vaccines. The vaccine components are unable to cross the placenta. The vaccine simply gives the body the recipe to make replicas of the virus protein, which in turn activate the immune response.

Vaccination primes the immune system with antibodies ready to block the COVID-19 from infecting our cells and specialist cells that can kill cells that do become infected. The second dose of the vaccine reinforces this immune response to produce higher levels of more effective and longer lasting protection.

COVID-19 vaccine safety in pregnancy

There are no safety concerns around giving Comirnaty to pregnant women/people or those planning pregnancy.

This vaccine is considered safe to use in pregnancy, based on two premises: firstly, that there is no known physiological mechanism by which the vaccine is likely to cause problems with pregnancy, and secondly, large-scale surveillance data do not indicate any safety concerns.

Although pregnant women were not formally included in the original clinical trials, the potential importance of immunising pregnant women against COVID-19 is now known, and several clinical studies are underway to improve the evidence around safe use of COVID-19 vaccines in pregnancy.^{18, 19}

To date, large-scale surveillance data have found no differences in pregnancy outcomes after vaccination with mRNA COVID-19 vaccines compared with the unvaccinated pregnant population.²⁰ The quantity of safety data in people who are pregnant continues to increase worldwide.

mRNA COVID-19 vaccine Pfizer/BioNTech [mRNA-CV]

For health professional use

In the US, surveillance data (v-Safe) showed the rate of miscarriage was consistent with the expected rate and did not increase following receipt of mRNA COVID-19 vaccines preconception or in early pregnancy (prior to 20 weeks' gestation).²¹ A larger Vaccine Safety Datalink study, involving over 100,000 pregnancies and 20,139 vaccinations, found that women who had experienced a miscarriage were not more likely to have been vaccinated within 28 days than those who had ongoing pregnancies.²²

When women who were vaccinated in pregnancy were compared with unvaccinated pregnant women, studies in the UK and Israel found no significant differences in any perinatal outcomes, including: stillbirth, fetal abnormalities postpartum haemorrhage, caesarean delivery, small for gestational age, maternal HDU or ICU admission, or preterm birth.^{23, 24}

Effectiveness of COVID-19 vaccination when given in pregnancy

COVID-19 vaccination in pregnancy is highly protective against severe disease, hospitalisation and death.

There is increasing international evidence that the effectiveness of Comirnaty when given in pregnancy is similar to that seen in the general population and that it is highly protective against symptomatic infection and hospitalisation after two doses.^{25, 26} The risk of COVID-19 infection was approximately five times less in those who were vaccinated in pregnancy than in unvaccinated pregnant people.²⁷ No fully vaccinated pregnant women were admitted to hospital for COVID-19 in the UK.¹⁰

This effectiveness is supported by the antibody response, vaccination was shown to induce a significantly higher protective antibody response than the natural viral infection.²⁸ There was no difference in antibody levels following vaccination between pregnant, lactating and non-pregnant women.^{28, 29}

Furthermore, there is mounting evidence of passive transfer of protective antibodies against COVID-19 from the mother to baby through the placenta.^{30, 31} Vaccination in the second or early third trimester has been suggested to be optimal for protective antibody transfer.³² Although, vaccination at any stage of pregnancy is recommended to protect the mother.

COVID-19 vaccination and breastfeeding

Breastfeeding women/people can receive a COVID-19 vaccine. There are no safety concerns associated with having this COVID-19 vaccine while breastfeeding or for the infant.

There is emerging evidence that vaccination in pregnancy or while breastfeeding provides temporary antibody protection to the baby through the cord blood and breastmilk.^{33, 34}

What are the likely responses to vaccination?

After hundreds of millions of doses of Comirnaty[®] given worldwide, the potential reactions to the vaccine have stayed consistent with those seen during the initial clinical trials. These include fatigue, headache, muscle aches, nausea, fever of 38–39°C and mild to moderate pain at the injection site. These reactions are more likely after the second dose vaccination.^{35, 36} Data from v-Safe found no differences in local and systemic responses between pregnant and non-pregnant women.²⁰

mRNA COVID-19 vaccine Pfizer/BioNTech [mRNA-CV]

For health professional use

Prior to receiving their vaccination, we recommend that pregnant women discuss the best ways to relieve possible post-vaccination discomfort and fever with their health professional. Paracetamol is considered safe during pregnancy to reduce headache or fever. Non-steroidal anti-inflammatories (NSAIDs) including ibuprofen and diclofenac should not be taken during pregnancy.

Anaphylaxis following vaccination is very rare (around five cases per million doses).³⁷ For this reason, all COVID-19 vaccine recipients are currently being asked to remain under observation for at least 15 minutes after receiving their vaccine. All vaccinators in New Zealand have training and equipment to manage anaphylaxis, should it occur.

Who should not receive a COVID-19 vaccine?

A COVID-19 vaccine is contraindicated (should not be given) for anyone who has had **anaphylaxis to an ingredient in the vaccine or a previous dose** of the same vaccine.

Who can receive a COVID-19 vaccine?

Women/people who are pregnant or who are planning a pregnancy can make an informed decision to receive Comirnaty™ (Pfizer/BioNTech) mRNA COVID-19 vaccine at any stage. Women planning pregnancy can receive mRNA COVID-19 vaccine at any time.

For anyone who is acutely unwell, fever >38°C or acute systemic illness, vaccination should be deferred until they are no longer acutely unwell.

References

1. Ministry of Health. COVID-19: Response planning: Ministry of Health; 2021 [updated 2021 October 26; ; cited 2021 November 01]. Available from: <https://www.health.govt.nz/our-work/diseases-and-conditions/covid-19-novel-coronavirus/covid-19-response-planning>
2. Principi N, Esposito S. Is the Immunization of Pregnant Women against COVID-19 Justified? *Vaccines* (Basel). 2021;9(9). doi: 10.3390/vaccines9090970
3. Zambrano LD, Ellington S, Strid P, et al. Update: Characteristics of Symptomatic Women of Reproductive Age with Laboratory-Confirmed SARS-CoV-2 Infection by Pregnancy Status - United States, January 22-October 3, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(44):1641-7. doi: 10.15585/mmwr.mm6944e3
4. Jamieson DJ, Theiler RN, Rasmussen SA. Emerging infections and pregnancy. *Emerg Infect Dis*. 2006;12(11):1638-43. doi: 10.3201/eid1211.060152
5. Branch DW. Physiologic adaptations of pregnancy. *Am J Reprod Immunol*. 1992;28(3-4):120-2. doi: 10.1111/j.1600-0897.1992.tb00771.x
6. Robinson DP, Klein SL. Pregnancy and pregnancy-associated hormones alter immune responses and disease pathogenesis. *Horm Behav*. 2012;62(3):263-71. doi: 10.1016/j.yhbeh.2012.02.023
7. Villar J, Ariff S, Gunier RB, et al. Maternal and Neonatal Morbidity and Mortality Among Pregnant Women With and Without COVID-19 Infection: The INTERCOVID Multinational Cohort Study. *JAMA Pediatr*. 2021. doi: 10.1001/jamapediatrics.2021.1050
8. Magnus MC, Oakley L, Gjessing HK, et al. Pregnancy and risk of COVID-19: a Norwegian registry-linkage study. *BJOG*. 2021 (preprint). doi: 10.1111/1471-0528.16969
9. Allotey J, Stallings E, Bonet M, et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ*. 2020;370:m3320. doi: 10.1136/bmj.m3320
10. Vousden N, Ramakrishnan R, Bunch K, et al. Impact of SARS-CoV-2 variant on the severity of maternal infection and perinatal outcomes: Data from the UK Obstetric Surveillance System national cohort. *medRxiv*. 2021 (preprint):2021.07.22.21261000. doi: 10.1101/2021.07.22.21261000

mRNA COVID-19 vaccine Pfizer/BioNTech [mRNA-CV]

For health professional use

11. Sun H. Approximation and evaluation of the spontaneous abortion rate with COVID-19 vaccination in pregnancy. *Am J Obstet Gynecol* MFM. 2021;100510. doi: 10.1016/j.ajogmf.2021.100510
12. Ghelichkhani S, Jenabi E, Jalili E, et al. Pregnancy outcomes among SARS-CoV-2-infected pregnant women with and without underlying diseases: a case-control study. *J Med Life*. 2021;14(4):518-22. doi: 10.25122/jml-2021-0157
13. Mullins E, Hudak ML, Banerjee J, et al. Pregnancy and neonatal outcomes of COVID-19: co-reporting of common outcomes from PAN-COVID and AAP SONPM registries. *Ultrasound Obstet Gynecol*. 2021. doi: 10.1002/uog.23619
14. Karasek D, Baer RJ, McLemore MR, et al. The association of COVID-19 infection in pregnancy with preterm birth: A retrospective cohort study in California. *Lancet Reg Health Am*. 2021;2:100027. doi: 10.1016/j.lana.2021.100027
15. Liguoro I, Pilotto C, Bonanni M, et al. SARS-CoV-2 infection in children and newborns: a systematic review. *European Journal of Pediatrics*. 2020;179(7):1029-46. doi: 10.1007/s00431-020-03684-7
16. Walker KF, O'Donoghue K, Grace N, et al. Maternal transmission of SARS-CoV-2 to the neonate, and possible routes for such transmission: a systematic review and critical analysis. *BJOG*. 2020;127(11):1324-36. doi: 10.1111/1471-0528.16362
17. Knezevic I, Liu MA, Peden K, et al. Development of mRNA Vaccines: Scientific and Regulatory Issues. *Vaccines (Basel)*. 2021;9(2). doi: 10.3390/vaccines9020081
18. Pfizer and BioNTech commence global clinical trial to evaluate COVID-19 vaccine in pregnant women [press release]. New York, US and Mainz, Germany: Pfizer, 2021 February 18 2021. 'Available from': <https://www.pfizer.com/news/press-release/press-release-detail/pfizer-and-biontech-commence-global-clinical-trial-evaluate>
19. Centers for Disease Control and Prevention. V-safe COVID-19 Vaccine Pregnancy Registry: US Dept of Health and Human Services; 2021 [updated 2021 June 2021 June 3]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/safety/vsafepregnancyregistry.html>
20. Shimabukuro TT, Kim SY, Myers TR, et al. Preliminary Findings of mRNA Covid-19 Vaccine Safety in Pregnant Persons. *N Engl J Med*. 2021;384(24):2273-82. doi: 10.1056/NEJMoa2104983
21. Zauche LH, Wallace B, Smoots AN, et al. Receipt of mRNA Covid-19 Vaccines and Risk of Spontaneous Abortion. *New England Journal of Medicine*. 2021;385(16):1533-5. doi: 10.1056/NEJMc2113891
22. Kharbanda EO, Haapala J, DeSilva M, et al. Spontaneous Abortion Following COVID-19 Vaccination During Pregnancy. *JAMA*. 2021. doi: 10.1001/jama.2021.15494
23. Blakeway H, Prasad S, Kalafat E, et al. COVID-19 vaccination during pregnancy: coverage and safety. *Am J Obstet Gynecol*. 2021. doi: 10.1016/j.ajog.2021.08.007
24. Wainstock T, Yoles I, Sergienko R, et al. Prenatal maternal COVID-19 vaccination and pregnancy outcomes. *Vaccine*. 2021;39(41):6037-40. doi: 10.1016/j.vaccine.2021.09.012
25. Dagan N, Barda N, Biron-Shental T, et al. Effectiveness of the BNT162b2 mRNA COVID-19 vaccine in pregnancy. *Nat Med*. 2021;27(10):1693-5. doi: 10.1038/s41591-021-01490-8
26. Bleicher I, Kadour-Peero E, Sagi-Dain L, et al. Early exploration of COVID-19 vaccination safety and effectiveness during pregnancy: interim descriptive data from a prospective observational study. *Vaccine*. 2021;39(44):6535-8. doi: 10.1016/j.vaccine.2021.09.043
27. Goldshtein I, Nevo D, Steinberg DM, et al. Association Between BNT162b2 Vaccination and Incidence of SARS-CoV-2 Infection in Pregnant Women. *JAMA*. 2021;326(8):728-35. doi: 10.1001/jama.2021.11035
28. Gray KJ, Bordt EA, Atyeo C, et al. Coronavirus disease 2019 vaccine response in pregnant and lactating women: a cohort study. *Am J Obstet Gynecol*. 2021;225(3):303 e1- e17. doi: 10.1016/j.ajog.2021.03.023
29. Collier AY, McMahan K, Yu J, et al. Immunogenicity of COVID-19 mRNA Vaccines in Pregnant and Lactating Women. *JAMA*. 2021. doi: 10.1001/jama.2021.7563
30. Prabhu M, Murphy EA, Sukhu AC, et al. Antibody Response to Coronavirus Disease 2019 (COVID-19) Messenger RNA Vaccination in Pregnant Women and Transplacental Passage Into Cord Blood. *Obstetrics & Gynecology*. 2021;138(2):278-80. doi: 10.1097/aog.0000000000004438
31. Beharier O, Plitman Mayo R, Raz T, et al. Efficient maternal to neonatal transfer of antibodies against SARS-CoV-2 and BNT162b2 mRNA COVID-19 vaccine. *J Clin Invest*. 2021;131(13). doi: 10.1172/jci150319
32. Gee S, Chandiramani M, Seow J, et al. The legacy of maternal SARS-CoV-2 infection on the immunology of the neonate. *Nat Immunol*. 2021. doi: 10.1038/s41590-021-01049-2
33. Golan Y, Prahli M, Cassidy AG, et al. COVID-19 mRNA Vaccination in Lactation: Assessment of adverse effects and transfer of anti-SARS-CoV2 antibodies from mother to child. *medRxiv*. 2021. doi: 10.1101/2021.03.09.21253241
34. Perl SH, Uzan-Yulzari A, Klainer H, et al. SARS-CoV-2-Specific Antibodies in Breast Milk After COVID-19 Vaccination of Breastfeeding Women. *JAMA*. 2021. doi: 10.1001/jama.2021.5782
35. Medsafe. Overview of Vaccine Reports 2021 [updated updated weekly; cited 2021 November 01]. Available from: <https://www.medsafe.govt.nz/COVID-19/vaccine-report-overview.asp>
36. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020;383(27):2603-15. doi: 10.1056/NEJMoa2034577
37. Shimabukuro TT, Cole M, Su JR. Reports of Anaphylaxis After Receipt of mRNA COVID-19 Vaccines in the US-December 14, 2020-January 18, 2021. *JAMA*. 2021. doi: 10.1001/jama.2021.1967